Author Index

Aomura, Y., see Ogino, H. 389
Ashraf, S. M.
——, Rajiah, A., Krishna, M. R. and Rao, M. B.
Limiting activity coefficients of aliphatic alcohols in phthalates 163
Astrua Testori, S., see Maisano, F. 422
Attygalle, A. B.
——, Zlatkis, A. and Middleditch, B. S.
Derivatization with 1,1-dimethylhydrazine for identification of carbonyl compounds resulting from ozonolysis 284
Bergenhem, N.
—— and Carlsson, U.
Some new by-products in primary structure analysis with dimethylaminoazobenzene isothiocyanate as an aid in the thin-layer chromatographic identification of residues 318
Bertolo, P. L., see Gennaro, M. C. 433
Betti, A., see Coppi, S. 406
Biacs, P., see Daood, H. G. 296
Billedeau, S. M.
Fluorimetric determination of vitamin K_3 (menadione sodium bisulfite) in synthetic animal feed by high-performance liquid chromatography using a post-column zinc reducer 371
Brown, S. Y., see Garland, W. A. 250
Brunner, C. A.
—— and Wainer, I.
Direct stereochemical resolution of enantiomeric amides via thin-layer chromatography on a covalently bonded chiral stationary phase 277
Budahegyi, M. V., see Tarján, G. 1
Cabras, P.
——, Spanedda, L., Pellecchia, M. and Gennari, M.
High-performance liquid chromatographic determination of the herbicide terbutylazine and its dealkylated metabolites in soil 411
Cabré, F.
——, Canela, E. I. and Canela, M. A.
Accuracy and precision in the determination of Stokes radii and molecular masses of proteins by gel filtration chromatography 347
Cañas, C., Igualda, see Mañes Vinuesa, J. 365
Canela, E. I., see Cabré, F. 347
Canela, M. A., see Cabré, F. 347
Carlsson, U., see Bergenhem, N. 318
Carr, R. J. G., see Jager, B. J. 331
Caude, M., see Lienne, M. 265
Chen, S.-T., see Lo, L.-C. 336
Cheynier, V.
——, Rigaud, J. and Moutounet, M.
High-performance liquid chromatographic determination of the free *o*-quinones of *trans*-caffeoyltartaric acid, 2-S-glutathionylcaffeoyltartaric acid and catechin in grape must 428
Collins, P., see Newsome, W. H. 416
Coppi, S.
—— and Betti, A.
Liquid chromatographic separation and behaviour of some triazines on styrene–divinylbenzene columns 406
Cortés, J. C., Moltó, see Mañes Vinuesa, J. 365
Crews, T., see Garland, W. A. 250
Cristalli, M., see Fanali, S. 441
Czinkotai, B., see Daood, H. G. 296
Daood, H. G.
——, Czinkotai, B., Hoschke, Á. and Biacs, P.
High-performance liquid chromatography of chlorophylls and carotènoids from vegetables 296
De Vos, R. H., see Heeremans, C. E. M. 219
Dreux, M., see Lafosse, M. 209
Elk, R., van, see Li, K. W. 445
Evans, M. B.
—— and Haken, J. K.
Recent developments in the gas chromatographic retention index scheme (Review) 93
Fanali, S.
——, Cristalli, M. and Quaglia, M. G.
Determination of rubidium, sodium, calcium and thiamine in a pharmaceutical preparation by capillary isotachophoresis 441
Fenelonov, V. B., see Zheivot, V. I. 155
Fields, S. M.
——, Grether, H.-J. and Grolimund, K.
Analysis of organic isocyanates using capillary supercritical-fluid chromatography 175
Fields, S. M.
—— and Grolimund, K.
Evaluation of supercritical sulfur hexafluoride as a mobile phase for polar and non-polar compounds 197
Font Pérez, G., see Mañes Vinuesa, J. 365
Fornstedt, N.
—— and Porath, J.
Sulphated Sepharose —a strong cation exchanger 261
Fukuda, E. K., see Garland, W. A. 250

Garland, W. A.
——, Crews, T., Brown, S. Y. and Fukuda, E. K.
Comparison of manual and exponentially modified gaussian based methods for the determination of the peak heights of selected-ion current profiles acquired in a mass spectral drug assay 250
Gavrilov, V. Yu., see Zheivot, V. I. 155
Gennari, M., see Cabras, P. 411
Gennaro, M. C.
—— and Bertolo, P. L.
Determination of the principal anionic components in wines and soft drinks, by ion interaction reversed-phase high-performance liquid chromatography 433
Geraerts, W. P. M., see Li, K. W. 445
Goward, C. R., see Jager, B. J. 331
Grandi, G., see Maisano, F. 422
Greef, J., van der, see Heeremans, C. E. M. 219
Greenaway, W.
——, May, J. and Whatley, F. R.
Flavonoid aglycones identified by gas chromatography–mass spectrometry in bud exudate of *Populus balsamifera* 393
Grether, H.-J., see Fields, S. M. 175
Grolimund, K., see Fields, S. M. 175, 197
Guan, Y.
——, Kiraly, J. and Rijks, J. A.
Interactive retention index database for compound identification in temperature-programmed capillary gas chromatography 129
Györ, M., see Tarján, G. 1
Haken, J. K., see Evans, M. B. 93
Hanaoka, R., see Sugii, A. 357
Harada, K., see Sugii, A. 357
Heeremans, C. E. M.
——, Van der Hoeven, R. A. M., Niessen, W. M. A., Vuik, J., De Vos, R. H. and Van der Greef, J.
Specific screening methods for glucosinolates in sprout extracts using on-line thermospray liquid chromatography–tandem mass spectrometry 219
Helliger, W., see Lindner, H. 243
Herbreteau, B., see Lafosse, M. 209
Hoeven, R. A. M., van der, see Heeremans, C. E. M. 219
Hoschkè, Á., see Daood, H. G. 296
Igualada Cañas, C., see Mañes Vinuesa, J. 365
Itabashi, Y.
——, Takagi, T. and Tsuda, T.
High-performance liquid chromatographic separation of alkane-1,2-diol enantiomers on a chiral slurry-packed capillary column 271
Jager, B. J.
——, Carr, R. J. G. and Goward, C. R.
Ultra-thin electrophoretic gel densitometry using photo-thermal deflection spectroscopy 331
Janda, V.
—— and Van Langenhove, H.
Determination of chlorophenols in water by direct acetylation and solid-phase extraction 327
Jelínek, I., see Snopek, J. 308
Joosse, J., see Li, K. W. 445
Kaneta, T., see Tanaka, S. 303
Kiraly, J., see Guan, Y. 129
Koza, T., see Řezanka, T. 290
Krishna, M. R., see Ashraf, S. M. 163
Kuš, P., see Śliwiok, J. 314
Kysilka, R., see Řezanka, T. 290
Lafosse, M.
——, Herbreteau, B., Dreux, M. and Morin-Allory, L.
Controle de certains systemes de chromatographie liquide haute performance à l'aide d'un détecteur évaporatif à diffusion de lumière 209
Langenhove, H., Van, see Janda, V. 327
Laurensen, J. J.
—— and Nouws, J. F. M.
Simultaneous determination of nitrofuran derivatives in various animal substrates by high-performance liquid chromatography 321
Li, K. W.
——, Geraerts, W. P. M., Van Elk, R. and Joosse, J.
Dot immunobinding assay of high-performance liquid chromatographic fractions on poly(vinylidene difluoride) membranes 445
Lienne, M.
——, Caude, M., Rosset, R. and Tambuté, A.
Direct resolution of anthelmintic drug enantiomers on Chiral-AGP protein-bonded chiral stationary phase 265
Lindner, H.
——, Wesierska-Gadek, J., Helliger, W., Puschendorf, B. and Sauermann, G.
Identification of ADP-ribosylated histones by the combined use of high-performance liquid chromatography and electrophoresis 243
Lo, L.-C.
——, Chen, S.-T., Wu, S.-H. and Wang, K.-T.
Separation of diastereomers of protected dipeptides by normal-phase high-performance liquid chromatography 336
Lombosi, E. R., see Tarján, G. 1
Lombosi, T. S., see Tarján, G. 1

Maisano, F.
——, Astrua Testori, S. and Grandi, G.
Immobilized metal-ion affinity chromatography of human growth hormone 422
Malakhov, V. V., see Zheivot, V. I. 155
Mañes Vinuesa, J.
——, Moltó Cortés, J. C., Igualada Cañas, C. and Font Pérez, G.
Isolation and concentration of organophosphorus pesticides from water using a C_{18} reversed phase 365
Masuda, S., see Sugii, A. 357
Matsumura, G., see Shimada, E. 145
May, J., see Greenaway, W. 393
Mészáros, S. Y., see Tarján, G. 1
Meuldijk, J., see Verhaar, L. A. Th. 381
Middleditch, B. S., see Attygalle, A. B. 284
Moltó Cortés, J. C., see Mañes Vinuesa, J. 365
Morin-Allory, L., see Lafosse, M. 209
Morrishow, A. M., see Prell, G. D. 256
Moutounet, M., see Cheynier, V. 428
Newsome, W. H.
—— and Collins, P.
Multiresidue method for fungicide residues in fruit and vegetables 416
Niessen, W. M. A., see Heeremans, C. E. M. 219
Nishimura, K., see Sugii, A. 357
Nouws, J. F. M., see Laurensen, J. J. 321
Nyiredy, Sz., see Tarján, G. 1
Ogino, H.
——, Suzuki, T. and Aomura, Y.
Simple modification to compensate for the baseline variation due to the introduction of a reduced-pressure gas sample into a gas chromatograph 389
Ovsyan-Nikova, I. A., see Zheivot, V. I. 155
Parkin, J. E.
High-performance liquid chromatography of mercury and phenylmercury as the N-disubstituted dithiocarbamate complexes 401
Pellecchia, M., see Cabras, P. 411
Pérez, G., Font, see Mañes Vinuesa, J. 365
Porath, J., see Fornstedt, N. 261
Prell, G. D.
—— and Morrishow, A. M.
Altered retention of derivatives of *tele*-methylimidazoleacetic acid in acid-hydrolyzed samples of brain measured by capillary gas chromatography and mass spectrometry 256
Puschendorf, B., see Lindner, H. 243
Quaglia, M. G., see Fanali, S. 441
Rajiah, A., see Ashraf, S. M. 163
Rao, M. B., see Ashraf, S. M. 163
Řezanka, T.
——, Koza, T., Kysilka, R., Víden, I. and Wurst, M.
Reversed-phase liquid chromatography for enrichment of very-long-chain fatty acids and their identification by gas chromatography–mass spectrometry 290
Rigaud, J., see Cheynier, V. 428
Rijks, J. A., see Guan, Y. 129
Rosset, R., see Lienne, M. 265
Sajewicz, M., see Śliwiok, J. 314
Sauermann, G., see Lindner, H. 243
Schmid, J. P., see Schwarzenbach, R. 231
Schwarzenbach, R.
—— and Schmid, J. P.
Determination of N-nitrosodiethanolamine in cosmetics. High-performance liquid chromatography and gas chromatography–mass spectrometry as alternative methods to chemiluminescence detection 231
Shalaeva, M. E., see Zheivot, V. I. 155
Shimada, E.
—— and Matsumura, G.
Determination of Stokes radii of hyaluronate oligosaccharides by Sephadex gel chromatography 145
Śliwiok, J.
——, Kuś, P. and Sajewicz, M.
Some data on the thin-layer and high-performance liquid chromatographic separation of some dithia[3.3]phanes 314
Smolková-Keulemansová, E., see Snopek, J. 308
Snopek, J.
——, Jelínek, I. and Smolková-Keulemansová, E.
Use of cyclodextrins in isotachophoresis. VIII. Two-dimensional chiral separation in isotachophoresis 308
Spanedda, L., see Cabras, P. 411
Sugii, A.
——, Harada, K., Nishimura, K., Hanaoka, R. and Masuda, S.
High-performance liquid chromatography of proteins on N-methylpyridinium polymer columns 357
Suzuki, T., see Ogino, H. 389
Takács, J. M., see Tarján, G. 1
Takagi, T., see Itabashi, Y. 271
Tambuté, A., see Lienne, M. 265
Tanaka, S.
——, Kaneta, T. and Yoshida, H.
Isotachophoretic separation of cyano complexes of noble metals using ion-pairing equilibrium in water–acetonitrile solvent 303

Tarján, G.
——, Nyiredy, Sz., Györ, M., Lombosi, E. R., Lombosi, T. S., Budahegyi, M. V., Mészáros, S. Y. and Takács, J. M.
Thirtieth anniversary of the retention index according to Kováts in gas–liquid chromatography (Review) 1
Testori, S., Astrua, see Maisano, F. 422
Tsuda, T., see Itabashi, Y. 271
Van der Greef, J., see Heeremans, C. E. M. 219
Van der Hoeven, R. A. M., see Heeremans, C. E. M. 219
Van der Wiele, K., see Verhaar, L. A. Th. 381
Van Elk, R., see Li, K. W. 445
Van Langenhove, H., see Janda, V. 327
Verhaar, L. A. Th.
——, Meuldijk, J. and Van der Wiele, K.
High-performance liquid chromatography of mixtures of catechol and its reaction products with glyoxylic acid 381
Víden, I., see Řezanka, T. 290
Vinuesa, J., Mañes, see Mañes Vinuesa, J. 365
Vos, R. H., de, see Heeremans, C. E. M. 219
Vuik, J., see Heeremans, C. E. M. 219
Wainer, I., see Brunner, C. A. 277
Wang, K.-T., see Lo, L.-C. 336
Wesierska-Gadek, J., see Lindner, H. 243
Whatley, F. R., see Greenaway, W. 393
Wiele, K., van der, see Verhaar, L. A. Th. 381
Wu, S.-H., see Lo, L.-C. 336
Wurst, M., see Řezanka, T., 290
Yoshida, H., see Tanaka, S. 303
Zheivot, V. I.
——, Shalaeva, M. E., Gavrilov, V. Yu., Fenelonov, V. B., Ovsyan-Nikova, I. A. and Malakhov, V. V.
Intermolecular interactions in the sorbate-modified graphitized thermal carbon black system 155
Zlatkis, A., see Attygalle, A. B. 284

Errata

J. Chromatogr., 465 (1989) 87–93
p. 92, Table I, first column, "μmol/l" should read "μmol/24-h".

J. Chromatogr., 467 (1989) 332–335
p. 333, 8th line of the *Validation* section, "concentration range 0.025 to 0.075 mg/ml" should read "0.05 to 0.15 mg/ml".
p. 335, ref. 1, the volume number is 227.